

4/15/2019

Briarcrest Veterinary Care Center
7313 E Tanque Verde Rd
Tucson, AZ 85715
(520) 298-4123

Payment On Account

Dane Haven Rescue
P O Box 27061
Prescott Valley, AZ 86312

Client ID: 1535

Previous Balance:	\$10.13
L/Visa, :	(\$10.13)
Subtotal:	<u>\$0.00</u>
Change Given:	
Balance Due:	<u><u>\$0.00</u></u>

Thank You!

Current

30 Days

60 Days

90 Days

\$0.00

\$0.00

\$0.00

\$0.00

Briarcrest Veterinary Care Center7313 E Tanque Verde Rd
Tucson, AZ 85715
(520) 298-4123

Page 1 / 1

Dane Haven Rescue
P O Box 27061
Prescott Valley, AZ 86312Client ID: 1535
Invoice #: 296584
Date: 4/12/2019

Patient ID: 1535-4	Species: CANINE	Weight: 79.80 pounds
Patient Name: Daisy	Breed: GREAT DANE MIX	Birthdate: 04/08/2015 Sex: Spayed Female

	<u>Description</u>	<u>Staff Name</u>	<u>Quantity</u>	<u>Total</u>
4/11/2019	Courtesy Medical Exam	Suzie Maki, DVM	1.00	\$0.00
	Fluoxetine 20mg		7.00	\$12.66
		Patient Subtotal:		\$12.66

Instructions

Your invoice was processed by rd

Invoice Total:	\$12.66
Preferred: 20:	(\$2.53)
Not Taxable :	\$0.00
Total:	\$10.13
Balance Due:	\$10.13
Previous Balance:	\$0.00
Balance Due:	\$10.13

Check out our new on-line store by visiting our website at My-Vet.com - Ordering food, flea & tick items along with other products is just a click away. In-home delivery, auto shipping and reminders are available. Ask any staff member for details!

Briarcrest Veterinary Care Center

520-298-4123

This medication has been prescribed for **Daisy Dane Haven Rescue**

Fluoxetine (for veterinary information only)

Brand Name: Reconcile, Prozac

Available in 10 mg tablets; 10 mg, 20 mg, 40 mg capsules; oral liquid; and 8 mg, 16 mg, 32 mg, 64 mg chewable tablets for dogs

Anxiety, obsessive-compulsive disorder, and depression are common problems for humans. Because the development of drugs that alleviate these conditions is ongoing, there are currently many medications on the market for these uses. One of the more popular classes of anti-anxiety medications are the selective serotonin reuptake inhibitors, called SSRIs for short.

Serotonin is a neurotransmitter in the brain associated, among other things, with mood elevation and reduced aggression. Increasing serotonin in the brain means less anxiety and a happier attitude. By inhibiting the brain's system for removing used serotonin, SSRIs cause serotonin to linger, lasting longer. The more serotonin we have in our brains, the less anxiety, obsession, and depression we get.

The pharmaceutical company Eli Lilly tested many related compounds and released fluoxetine as the most specific SSRI of all of them. Fluoxetine, marketed under the brand name Prozac®, was not quite the first SSRI on the human market but quickly became the most popular.

It was not long after that that fluoxetine found its way into veterinary use for animals with anxiety, compulsive behavior, and other behavior issues. When low-cost generic fluoxetine became available, its use increased in the veterinary market and in 2007 Eli Lilly released a version of fluoxetine specifically labeled for animal use.

How this Medication is Used

Fluoxetine has been used in many animal behavior issues: inappropriate urine marking in dogs and cats; separation anxiety; compulsive chewing; circling; self-mutilation; and even aggression. This product is specifically approved for the treatment of separation anxiety. It should be noted that the treatment of behavior disorders requires training in addition to simply giving the pet a medication. Fluoxetine is usually given once daily.

Side Effects

It is important to understand that whenever an anti-anxiety medication is used, the phenomenon of disinhibition is possible. What this means is that an animal's inhibitions about aggressive behavior may be reduced when his or her anxiety over the consequences of such behavior is removed. An animal that was not previously aggressive could potentially become aggressive.

The most common side effect is drowsiness or lethargy. Appetite reduction is a common side effect of fluoxetine in dogs. Some patients will experience an upset stomach with this medication.

Interactions with Other Drugs

Fluoxetine should not be used in combination with drugs that could increase the likelihood of seizures, such as acepromazine. Fluoxetine should not be given in conjunction with a monoamine oxidase inhibitor (MAO inhibitor) such as selegiline or amitraz. This includes amitraz-containing collars such as the Preventic® collar, dips such as Mitaban® dip, and flea products such as Promeris® for dogs.

Concerns and Cautions

This medication is best not used in patients with diabetes mellitus or with seizure disorders.

Patient: Daisy	Owner: Dane Haven Rescue	Chart: 1535	Date: 4/11/2019
----------------	--------------------------	-------------	-----------------

History/Presenting Complaint: 4 Yrs. 0 Mos. GREAT DANE MIX Spayed Female - Initial Exam. Also having issues with submissive urination. Started about a week ago.

Meds: none
CR

<input type="checkbox"/> Fractious	<input type="checkbox"/> Brief Exam	<input type="checkbox"/> No Exam	Weight (lbs)	Diet
<input type="checkbox"/> Difficult	<input type="checkbox"/> Problem Specific	<input type="checkbox"/> Comprehensive	79.8	Blue Buffalo High Protein

General Appearance <input checked="" type="checkbox"/> NRM <input type="checkbox"/> ABM <input type="checkbox"/> NE nervous, tail tucked, wide eyes, lip licking.	Oral Cavity <input checked="" type="checkbox"/> NRM <input type="checkbox"/> ABM <input type="checkbox"/> NE	Teeth & Gingiva <input checked="" type="checkbox"/> NRM <input type="checkbox"/> ABM <input type="checkbox"/> NE minimal calculus	Eyes <input checked="" type="checkbox"/> NRM <input type="checkbox"/> ABM <input type="checkbox"/> NE
Ears <input checked="" type="checkbox"/> NRM <input type="checkbox"/> ABM <input type="checkbox"/> NE	Heart <input checked="" type="checkbox"/> NRM <input type="checkbox"/> ABM <input type="checkbox"/> NE	Respiratory <input checked="" type="checkbox"/> NRM <input type="checkbox"/> ABM <input type="checkbox"/> NE	Abdomen <input checked="" type="checkbox"/> NRM <input type="checkbox"/> ABM <input type="checkbox"/> NE
Musculoskeletal <input type="checkbox"/> NRM <input checked="" type="checkbox"/> ABM <input type="checkbox"/> NE guards hind - unable to examine hips ambulates normally in room.	Lymph Nodes <input checked="" type="checkbox"/> NRM <input type="checkbox"/> ABM <input type="checkbox"/> NE	Genitourinary <input checked="" type="checkbox"/> NRM <input type="checkbox"/> ABM <input type="checkbox"/> NE normal vulvar conformation, no perivulvar dermatitis.	Integument <input checked="" type="checkbox"/> NRM <input type="checkbox"/> ABM <input type="checkbox"/> NE several scab old from fights with previous dog in home. left front near dewclaw, raised, bumpy white crusted dermal lesion with not well defined SQ swelling 8mm. - old puncture, old traumatized dermal mass?

Temp: 98.3ax	Pulse/HR: 90	Resp: 30	MM: Pink	CRT: 1-2 sec	Hydration: Normal
--------------	--------------	----------	----------	--------------	-------------------

BP: /	I.O.P. O.S. O.D.	Fecal Float: <input type="checkbox"/> POS <input type="checkbox"/> NEG	4DX: <input type="checkbox"/> POS <input type="checkbox"/> NEG	Clot
		Fecal Direct: <input type="checkbox"/> POS <input type="checkbox"/> NEG	FELV FIV: <input type="checkbox"/> POS <input type="checkbox"/> NEG	PCV TP

Pain Assesment Scale: 0 Body Condition Score: 3/5

Remarks:

A: Dx: Anxiety/phobia of men. Foster reports cowers, tucks tail, urinates, shows tummy when fiance (who is a man) gives her commands. She rolls into him for love and seeks his attention but behavior is only noted when talks to her. Does not occur with female foster. Recommend he drops treats for her without looking or interacting with her for a few days, since interacts well with him outside of speaking then give pets and treats. Feel will take time and anti-anxiety meds for her to come around. Also recommend behavioral consult/modification with Dr. Vanya Moreno. Licks paws incessantly at night. No head shaking, generalized pruritis. Suspect environmental allergy. Recommend OTC anti-histamine.

P: Recommend start Loratidine for paw licking if does not improve with Fluoxetine.

Quantity	Item Description	Directions	Refill Number
7.00	Fluoxetine 20mg	GIVE BY MOUTH 1 TABLET EVERY 24 HOURS FOR ONE WEEK, THEN GO TO THE 40MG DOSE THEREAFTER. AL-6	0

After one week time increase to 40mg dose. Written Rx for Fluoxetine 40mg PO SID # 30 - discussed possible side effects.

Recommend consult with Dr. Vanya Moreno.

Patient: Daisy	Owner: Dane Haven Rescue	Chart: 1535	Date: 4/11/2019
----------------	--------------------------	-------------	-----------------

PR call 2 weeks.

Dr. Suzie Maki

4/11/2019 ex sheet 1 Maki, DVM, Suzie
--